

Data Sheet

Catalogue Number	Protein Receptor Type IA Human Recombinant IY-380
<u> </u>	BMPR-1A, BMP-R1A, BMPR1A, BMR1A, CD292, CD-292, Serine/threonine-
Synonyms	protein kinase receptor R5, SKR5, Activin receptor-like kinase 3, ALK-3,
	ACVRLK3, EC 2.7.11.30, CD292 antigen.
Introduction	The bone morphogenetic protein (BMP) receptors are a family of
	transmembrane serine/threonine kinases that include the type I receptors
	BMPR1A and BMPR1B and the type II receptor BMPR2. These receptors
	are also closely related to the activin receptors, ACVR1 and ACVR2. The
	ligands of these receptors are members of the TGF-beta superfamily. TGF-
	betas and activins transduce their signals through the formation of
	heteromeric complexes with 2 different types of serine (threonine) kinase
	receptors: type I receptors of about 50-55 kD and type II receptors of
	about 70-80 kD. Type II receptors bind ligands in the absence of type I
	receptors, but they require their respective type I receptors for signaling,
	whereas type I receptors require their respective type II receptors for
	ligand binding.
Patent Rights	The sale and/or commercial use of Recombinant Adiponectin is prohibited
	in the United States of America (U.S.A).
	BMPR1A Human Recombinant extracellular domain produced in
	baculovirus is a monomeric, glycosylated, Polypeptide chain fused with
Description	6xHis tag at C-terminus and having a molecular mass of 23 kDa. The
	BMR1A is purified by proprietary chromatographic techniques.
Source	Insect Cells.
Physical Appearence	Sterile Filtered White lyophilized (freeze-dried) powder.
Thysical Appearence	CD292 was lyophilized from a concentrated (1mg/ml) sterile solution
Formulation	containing 1X PBS.
	It is recommended to reconstitute the lyophilized ALK-3 in sterile PBS not
Solubility	less than 100 μg/ml, which can then be further diluted to other aqueous
	solutions.
Stability	Lyophilized Bone Morphogenetic Protein Receptor 1A although stable at
	room temperature for 3 weeks, should be stored desiccated below
	-18°C. Upon reconstitution BMPR1A should be stored at 4°C between 2-7
	days and for future use below -18°C.
	For long term storage it is recommended to add a carrier protein (0.1%
	HSA or BSA).
	Please prevent freeze-thaw cycles.
Purity	Greater than 90.0% as determined by:
	(a) Analysis by RP-HPLC.
	(b) Analysis by SDS-PAGE.
Biological Activity	Measured by its ability to inhibit recombinant human BMP-2 induced
	alkaline phosphatase production by C2C12 myogenic cells. The ED50 for
	this effect is typically 1-3 µg/ml in the presence of 500 ng/ml of
	recombinant human BMP-2 corresponding to a Specific Activity of 2,000
	units/mg.

	Products are furnished for LABORATORY RESEARCH USE ONLY. The
Usage	product may not be used as drugs, agricultural or pesticidal products, food
	additives or household chemicals.